

Complementary Feeding Practices among Mothers of Children under Five Years of Age in Satellite Town, Lagos, Nigeria

Olatona F. A.^{1,*}, Odozi M. A.¹, Amu E. O.²

¹Department of Community Health and Primary Care, College of Medicine, University of Lagos, Nigeria
²Department of Epidemiology and Community Health, Ekiti State University, Ado-Ekiti, Ekiti State, Nigeria

Abstract In Nigeria, only 30 percent of infants ages 6-23 months are fed in accordance with the recommendation for infant and young children feeding practices despite the fact that it has been shown that complementary feeding is one of the most important preventive measures that would reduce the excess mortality for children under the age of five years. This study determined the complementary feeding practices, among mothers of children under the age of five years in satellite town, Lagos. It was a descriptive cross sectional study which employed a multistage sampling method to select 358 mothers whose last confinement was not more than five years to prevent recall bias. Data was collected using structured interviewer-administered questionnaires and analyzed using Epi-Info version 6 computer software. Most of the respondents (65.0%, n= 227) have completed tertiary while only 1.2 % (n = 3) had no formal education. Less than half of the respondents (48.4%) introduced drinks or foods at age 6months. Even though majority (91.9%) continued breast feeding after introducing other foods, more than half (57.1%) discontinued it before 12 months. Twenty four hours dietary recall revealed that the commonest food given was cereals (65.9%) followed by fruits (46.4%) and tubers (42.7%). Most of the respondents fed the child responsively (70.3%) and prepared foods hygienically (75.1%). Complementary feeding practices though better than in many other communities within and outside Nigeria were inadequate compared to the recommended standards. Complementary feeding education using various media especially health facilities is very important for mothers of infants.

Keywords Complementary feeding, practices, mothers of under-five, Nigeria

1. Introduction

In Nigeria, under-nutrition remains a public health problem. According to the 2008 National Demographic and Health Survey, 23% of children are underweight, 14% are wasted and 41% are stunted [1]. The development of children's full human potential depends on adequate nutrition during infancy and early childhood. Exclusive breastfeeding is adequate for the first six months of life but timely introduction of nutritionally-adequate, safe, age-appropriate complementary feeding at sixth month of age is recommended for better health and development of infants [2]. The World Health Organization recommends that breastfeeding should continue with appropriate complementary feeding from 6 months to 2 years and beyond [3].

According to the Nigeria Demographic and Health Survey 2008, only 30 percent of infants aged 6-23 months and living

are fed in accordance with the recommendation for infant and young children feeding practices [1]. Problems associated with complementary feeding in particular include poorly timed introduction of complementary foods (too early or too late), bottle-feeding, infrequent feeding, unsupervised feeding, poor feeding methods, poor hygiene, and child care practices [4, 5].

Malnutrition is responsible for more than 41% of the deaths that occur annually in children from 6 to 24 months of age in developing countries which total approximately 2.3 million [6]. Moreover, one third of deaths among children under five years of age are attributable to nutrition and well above two thirds of these deaths, often associated with inappropriate feeding practices, occur during the first year of life [6]. Inadequate feeding of girls affects nutrient stores, leading to poor obstetrics outcome when they are having children later in life. Other consequences include growth failure and stunting, delayed sexual development, reduced muscle mass and strength, impaired intellectual development, weakened immune system, poor cognitive performance and learning achievement and increased lifetime risk of osteoporosis [7]. Every year, more than 200 million children under five years old fail to reach their full cognitive and

* Corresponding author:

folaton@gmail.com (Olatona F. A.)

Published online at <http://journal.sapub.org/fph>

Copyright © 2014 Scientific & Academic Publishing. All Rights Reserved

social potential [6]. Most of these children live in South Asia and sub-Saharan Africa [6].

Among preventive measures that would reduce the excess mortality for children under the age of five years, exclusive breast feeding and good quality complementary feeding have been listed as first and third, respectively, with a calculated 600,000 deaths per year preventable by good complementary feeding (i.e. 6% of deaths) [8]. Growth faltering occurs mainly in the first two years of life because of high nutritional needs for growth and development between 6-24 months [9]. In fact, the percentage of children who are underweight almost doubles from 14% among children less than 6 months of age to 26% among children age 12-17 because complementary foods are introduced to children in the latter group, thus increasing exposure to infections and susceptibility to illness [1].

In Delta State, Nigeria, a little above half (54.0%) of nursing mothers feed their infants with complementary foods three times daily, and 2.4% and 37.3% feed their infants using their hand and feeding bottle respectively.[10] In Osun State, many women (48.8%) terminate breast feeding because they feel children were old enough at age 12 months. Children are fed only when they are hungry (49.9%) and 89.2% used feeding bottle for feeding [11]. In some parts of Nigeria, complementary foods are introduced to majority of children much earlier at third month (41.2%) and some others at 1-2 months (17.8%) compared with the 6th month recommended [12].

However, little is known about complementary feeding practices among urban dwellers in Lagos, Nigeria. This study determined the complementary feeding practices among mothers of under-five in an urban area of Lagos State. The result will be useful for health educators to identify areas of focus for health education when educating the masses and policy makers in establishment and implementation of effective policies.

2. Methods

A cross-sectional descriptive study was conducted among mothers of children under-five years of age in Satellite town,

Lagos. The minimum sample size calculated using prevalence of appropriate infant and young child feeding practices in Nigeria, (30%) was 332 [1], however 358 mothers were selected for the study to make up for attrition and improperly completed questionnaire, using multi-stage sampling technique. In Satellite town there are seventy five (75) closes, fifteen (15) company owned estates, ten (10) privately owned estates and fifty (50) streets. Using the systematic sampling method with a sampling interval of five (5), every 5th close was selected out of the seventy five (75) closes. Fifteen (15) closes were therefore selected. Balloting was done to pick one between the two ends of each close to determine the starting point. One mother was interviewed from each household. All eligible mothers in the households in each close were interviewed until six (6) mothers were interviewed in each close to obtain ninety (90) respondents from the closes.

Two (2) streets were selected out of the fifty (50) streets using the balloting method. The average number of houses on each street was one hundred (100). Using the systematic random sampling with a sampling interval of two (2), every second (2nd) house was selected on each street to get fifty (50) houses. One (1) household was selected from each house using balloting method, to obtain fifty respondents from each street and therefore a total of one hundred respondents (100) from the streets section.

One (1) estate each from the company owned and privately owned estate was selected through balloting. Two (2) estates were therefore selected. The households in the estates did not have a systematic arrangement so all eligible mothers were interviewed until eighty five (85) mothers were interviewed in each estate to obtain one hundred and seventy (170) respondents from the estate section. A mother and child were included in the study if the youngest child was at least one year and less than 5 years of age. The youngest child was taken as the index child and all the responses obtained were based on this child. If more than one mother who met the inclusion criteria was seen in a household, one (1) mother was selected by the balloting technique.

Table 1. Breastfeeding and Complementary Feeding Practices for Index Child of Respondents

Breastfeeding and complementary feeding practices	Frequency	Percentage (%)
Child was breastfed	332	95.1
Continued breastfeeding after introducing other foods	305	91.9
Age at which drinks or foods were introduced (n=332)		
< 6 months	89	26.8
6 months	161	48.4
>6 months	82	24.8
Age when breastfeeding was discontinued (n=332)		
< 12 months	93	28.0
12 months	93	28.0
> 12 months	146	44.0

Table 2. Respondents' Pattern of Complementary Feeding of the Index Child

Feeding pattern (n=332)	Freq.	Percentage (%)
Responsive feeding (anytime the child gives the cue)	233	70.3
Correct frequency of foods at age 6 – 8 months (2 -3 times)	124	37.3
Correct frequency of foods at age 9 – 11 months (3 - 4times)	165	49.7
Correct frequency of foods at 12 months and above(4 - 6 times)	170	51.2

Table 3. Twenty Four (24) Hours Dietary Recall for Index Child of Respondents

All liquid or solid foods consumed	Frequency (n=349)	Percentage (%)
Plain water	313	87.4
Fresh fruits juice/fruits	166	46.4
Fresh animal milk/tea/coffee	143	39.9
Breast milk	100	27.9
Infant formula	85	23.7
Pap, pasta, rice, bread, biscuits.	236	65.9
Yam, garri, pounded yam, potatoes	153	42.7
Meat, chicken, fish or eggs	142	39.7
Vegetables	123	34.4
Beans, groundnut green peas	97	27.1

Table 4. Frequency of Total Food Consumption (Breast Milk or Complementary Food) In 24 Hours, Methods of Feeding Child and Hygiene Practices

Method of feeding and hygiene practices	Frequency (n=349)	Percentage (%)
Child was fed 7 times or more (breast milk & other foods)	37	10.5
Food thick enough to stay on a spoon	176	50.3
Thickness of food increased as child grew older	296	84.9
Chew food before giving to child	122	34.9
Mother feeds child	222	63.6
Used plate and spoon to feed child	280	80.2
Used feeding bottle to feed child	65	18.6
Hand feeding	4	1.2
Always washed hands before feeding child	281	80.6
Always sterilized feeding utensils	262	75.1

Data was collected in May 2013, using a pre-tested structured interviewer administered questionnaire. The questionnaire contained questions on complementary feeding practices. Four interviewers (secondary school leavers) were trained on how to administer the questionnaires correctly. If the interviewers didn't meet an eligible woman in the house at the time of visit, then they went back at a later time in the evening or at a later date. The completed questionnaires were inputted and analysed using Epi Info Statistical software (version 3.5.1).

Ethical approval was obtained from the Health Research and Ethics Committee of Lagos University Teaching Hospital and permission to conduct the research was obtained from the Local Council Development Area. Informed written consent was obtained from each respondent before administering the questionnaire. Utmost confidentiality of information obtained was ensured.

3. Results and Discussion

Most of the respondents (65.0%, n= 227) have completed tertiary while only 1.2% (n = 3) had no formal education. Almost all the respondents (95.1%) breastfed their children. This is similar to the findings of a study conducted in

Mauritius in which 93.4% of the respondents breast fed their babies [13]. This probably means that most of the mothers in this community have embraced baby friendly initiative. Almost half of the respondents (48.4%) introduced drinks or foods at age 6 months. The proportion who introduced complementary feeding at the age of 6 months is lower than that reported among mothers of under two children in Kassena-Nanka region of Ghana (61.2%) but slightly higher than the 37.6% reported in Brong Ahafo region in Ghana and 33.5% reported in South-West Nigeria [11, 14, 15]. It however contrasts with that reported in India where only 13% of the respondents had introduced complementary feeds at 6 months [16]. The higher responses reported in this and the Kassena-Nanka region of Ghana studies can be explained by the higher educational levels of the respondents and better access to information than those in the other studies.

Only about a quarter (26.8%) of the respondents in this study introduced complementary foods before the age of 6 months. This contrasts with the results obtained from some studies conducted in Gaza Strip, South-South, North-West and South-West regions of Nigeria where 79.9, 79.0 and 80.1 and 53.0% respectively introduced complementary feeding before the expected 6 months of age [10, 11, 12, 17]. The mothers in this study were probably more exposed to infant

feeding education and knew the danger of introducing meals before six months of age. Introduction of feeds before 6 months is fraught with the dangers of infections, allergies, digestive problems and obesity later in life and therefore should be discouraged [7, 18].

About a quarter, (24.8%) of the respondents equally introduced complementary foods after the age of 6 months which contrasts with that of a study conducted in India in which 63% of the respondents introduced complementary foods after 6 months [16]. After six months, breast milk is no longer sufficient to meet the nutritional requirements of an infant. Late introduction of complementary feeds leads to growth faltering which can predispose to overt malnutrition especially when associated with recurrent infections which are common about this stage [9].

A large percentage (91.9%) continued breast feeding after introducing other foods but well over half (56.1%) of them equally discontinued before or at 12 months; while only 43.9% discontinued after 12 months. The Kassena-Nanka Ghanaian study reported similar findings with 96.3% continuing breast feeding after introducing complementary feeds [14]. It is recommended that breast feeding should continue after introducing complementary feeds until child is two years old. This will give the necessary nutritional support while the child gradually gets accustomed to eating solid food [19].

Most of the respondents (70.3%) fed their children responsively while others fed their children at scheduled times or during family meal times. Responsive feeding is advocated by WHO and UNICEF as the best approach to ensure that infants are appropriately fed and has been shown to be associated with better mental development compared to traditional complementary feeding [7, 20].

As convenient as scheduled meal times may be, they might not coincide with the period of hunger in the infants, such that they might either refuse to eat, or take less or more than what they would ordinarily have eaten. Baby-led approach in complementary feeding may encourage greater satiety-responsiveness and healthy weight-gain in infants [21].

The World Health Organization (WHO) recommends certain age-specific frequencies of complementary feeding based on theoretical estimates of the energy provided by complementary foods. Infants should be fed initially 2-3 times a day between 6-8 months, increasing to 3-4 times daily between 9-11 months and 12-24 months with additional nutritious snacks offered 1-2 times per day, as desired [1]. In this study, only 37.3, 49.7 and 51.2% of the infants were fed according to these requirements at age 6-8, 9-11 and 12-24 months respectively.

Some studies conducted in Nigeria, India and Ghana similarly reported that the mothers did not know the appropriate age-specific frequency of feeding with complementary foods as stipulated by the WHO [10, 15, 16]. Most of them just fed their children as frequently as it seemed appropriate to them irrespective of the ages of the infants. This underscores the need for regular nutrition

education at various contact points where the mothers can be reached. Feeding less than the required number of times can lead to under-nutrition while feeding more than the recommended number of times can lead to over nutrition and later on obesity.

The twenty four hours (24 hours) dietary recall revealed that majority of the respondents gave cereal (65.9%) followed by fruits (46.4%) and then tubers (42.7%). Animal proteins in form of (meat fish or eggs) and vegetables were given by few people. A study conducted among children in Northern Uganda similarly reported that majority of the respondents (35.2%) were given cereals in the last 24 hours. However, the proportion that gave fruits and vegetables (31.1%) and beans (22.4%) was lower. The proportion that gave meat and milk /milk products were even much lower than that of the present study (1.8 and 0.5%) respectively. [22] Cereal based foods alone are not sufficient; in order to prevent micronutrient deficiencies, milk, meat products, legumes, fruits and vegetables should be integral parts of the complementary foods of children. Breast feeding should also be continued up till two years in order to further support their growth [6].

Only 10.5% of respondents fed their children more than 7 times as expected. This may be as a result of ignorance about this recommendation or because the parents could not afford to feed the infants more than seven times per day. This is similar to the report of a study conducted among returnee children in Northern Uganda in which majority of the children were fed only twice irrespective of their ages, in the 24 hours preceding the survey [22].

Majority of the children were fed by the mothers (63.6%). Mothers are better at practicing responsive feeding than other care givers some of which might be the older siblings. In responsive feeding the mother responds to the child's hunger and satiety cues, interacts with the child, experiments with different food combinations, tastes, textures, and methods of encouragement in case the child refuses many foods, and minimizes distractions during feeding. All these skills help to ensure that the child takes enough especially when ill [8, 23].

Only half of the respondents (50.3%) made the complementary food to be of the right consistency though majority (84.9%) increased the thickness as the children grew older. In the Brong Ahafo region study in Ghana however, only about one-third of caregivers reportedly gave thick, nutrient dense foods to children after 6 months of age while only 33.2% of them increased the thickness as the children grew older [15]. It is recommended that at the beginning of complementary feeding, foods should be semi-solid and soft (in the form of a puree) and should be crushed or mashed, never sifted or blended [24]. The consistency should then be gradually improved, considering the infant's eating skills and age. It should be noted that improper food consistency compromises the appropriate intake of nutrients by the infant [3].

About four-fifth of the respondents always washed their hands before feeding their children while three-quarters of

them always sterilized their feeding utensils. This contrasts with the report of a study conducted in Northern Nigeria in which only 28% of the respondents always washed their hands before feeding the children and only 19% always sterilized the feeding utensils [12]. The respondents in this study were more hygienic compared with their Northern counterparts. The fact that most of the mothers had tertiary education, higher socio-economic statuses and better access to amenities such as water and electricity compared to their counterparts in the North can explain the difference. It is expected that clean utensils be used to feed children in order to avoid introducing dirt and germs that might cause diarrhoea and other infections [24].

4. Conclusions

Less than half of the respondents (48.4%) introduced other drinks or foods at age 6 months. Even though majority (91.9%) continued breast feeding after introducing other foods, more than half (57.1%) discontinued it before 12 months. Complementary feeding practices in this study though better than in many other communities within and outside Nigeria were inadequate compared to the recommended standards. WHO guidelines on complementary feeding, should be communicated using various media to mothers of infants. Moreover, government should encourage female education so that more mothers will be educated and empowered to practice adequate and safe complementary feeding.

Conflict of Interest

We, the authors of this manuscript declare that there is no conflict of interest that might bias our work or inappropriately influence our judgment.

REFERENCES

- [1] National Population Commission, 2008a, 2007 Nigeria demographic and health survey, MD USA, NPC and ORC Macro Calverton, p.163-174.
- [2] World Health Organization, 2003, *Global strategy for infant and young child feeding*, Geneva, World Health Organization, p. 4-14.
- [3] Zere, E., McIntyre, D., 2003, Inequities in under five child malnutrition in South Africa, *International Journal for Equity in Health*, 2003, 11(2), 7.
- [4] Administrative Committee on Coordination/Subcommittee on Nutrition, 2010, Sixth Report on the World Nutrition Situation. Geneva.
- [5] Shordt, K., Cairncross, S., Sustainability of hygiene behaviour and the effectiveness of change interventions, London School of Hygiene and Tropical Medicine Research Online. 2004: Pg 34-48. Available from www.researchonline.lshm.ac.uk. (accessed 1st March 2013).
- [6] Sandoval-Priego, A.A., Reyes-Morals, H., Perez-Cuevas, D., Abrego-Blass, R., Orrico-Torres, E.S., 2003, Family strategies of life associated with malnutrition in children less than 2 years of age, *Salud Publica de Mex.*, 44, 1-9.
- [7] World Health Organization, 2003, Complementary feeding: report of the global consultation. Summary of guiding principles for complementary feeding of the breastfed child. WHO, Geneva. Available at: http://www.who.int/nutrition/topics/complementary_feeding/en/index.html (accessed 20 May, 2013)
- [8] Jones, G., Steketee, R.W., Black, R.E., Bhutta, Z.A., Morris, S.S., 2003, How many child deaths can we prevent this year? *Lancet*, 362 (9377), 65-71.
- [9] Institute of Medicine, 2006, Dietary reference intakes: the essential guide to nutrient requirements, National Academics Press, Washington D.C.
- [10] Osie-Efetie, B., Oyibo, P.G., Okperi, B.O., 2011, Weaning practices among nursing mothers in Ethiope East Local Government Area of Delta State, Nigeria. *Continental J. Biomed. Sci.*, 5(2), 19 – 28.
- [11] Ogunba, B.O., 2006, Maternal behavioural feeding practices and under-five nutrition: implication for child development and care. *J. Appl. Sci. Res.*, 2(12), 1132-1136.
- [12] Anigo, K.M., Ameh, D.A., Ibrahim, S., Danbauchi, S.S., 2009, Infant feeding practices and nutritional status of children in North Western Nigeria. *Asian J. Clin. Nutr.*, 1, 12-22.
- [13] Motee, A., Ramasawmy D., Pugo-Gunsam, P., Jeewon R., 2013, An assessment of the breast feeding and infant feeding pattern among mothers in Mauritius. *J. Nutr. Metabolism*. Available at: <http://dx.otg/10.1155/2013/243852> (accessed 10th June, 2013).
- [14] Adokiya M.N., 2010, Complementary feeding practices and nutrition status of young children 6-23 months of age in the Kassena-Nankana District, Upper East Region, Ghana. *Knust Space*. Available at <http://dSPACE.knust.edu.gh:8080/jspui/bitstream/123456789/103>. (accessed 12th March, 2014).
- [15] USAID, GHS, GAIN, PATHS and IYCN. Ghana promotion of complementary feeding practices project baseline survey report. December 2011. Page 26. Available at http://iycn.wpengine.netdna-cdn.com/files/IYCN_Ghana_Baseline_Report_1211. (accessed 5th February, 2014).
- [16] Kumudha, A., Khan M.E., Avishek, H., 2010, Increasing appropriate complementary feeding in rural Uttar Pradesh, India, *J. Fam. Welfare*, 56, 51-56.
- [17] Basil, J.K., Moain, E., Adnan, A., Abdel, M.L., Ayoub, A., 2011, Breast feeding, complementary feeding and weaning practices, among children up to 2 years old in Gaza Strip. *Ann. Alquds Med.*, 7, 15-26.
- [18] The Open University, Nutrition HEAT Module, Section 4, Infant and young child feeding. Available at: <http://labspace.open.ac.uk/mod/oucontent/view.php>. (accessed 6th April 2013).
- [19] Ashworth, A., Chopra, M., McCoy, D., Sanders, D., Jackson, D., Karaolis, N., et al., 2004, World Health Organization guidelines for management of severe malnutrition in rural

- South African hospitals: effect on case fatality and the influence of operational factors, *Lancet*, 363, 1110-1115.
- [20] Vazir, S., Engle, P., Balakrishna, N., Griffiths, P.L., Johnson S.L., Creed-Kanashiro, H., et al., 2013, Cluster-randomized trial on complementary and responsive feeding education to caregivers found improved dietary intake, growth and development among rural Indian toddlers. *Matern. Child Nutr.* 9(1), 99-117. doi: 10.1111/j.1740-8709.2012.00413
- [21] Brown, A., Lee, M.D., 2013, Early influences on child satiety-responsiveness: the role of weaning style. *Pediatr. Obes.*, doi: 10.1111/j.2047-6310.2013.00207.
- [22] Mokeri, A., Orikushaba, P., 2012, Nutritional status, complementary feeding practices and feasible strategies to promote nutrition in returnee children aged 6-23 months in Northern Uganda. *S Afr J Clin Nutr.*, 25(4), 173-179.
- [23] Similac.org. Your baby's growing digestive system. Available at: <http://similac.com/feeding-nutrition/baby-digestive-system>. (accessed 4th March 2013).
- [24] Kikafunda, J.K., Walker, A.F., Tumwine, J.K., 2003, Weaning foods and practices in Central Uganda: a cross-sectional study. *AJFAND* 3(2). [ISSN 1684-5378] Available at: <http://www.ajol.info/index.php/ajfand/issue/view/2491>. (accessed 26th March 2014).