

**BACTERIAL DIVERSITY STUDIES AND PROBIOTIC FUNCTIONS OF
BACILLUS SPECIES IN FERMENTED LOCUST BEAN SEEDS (PARKIA
BIGLOBOSA JACQ. BENTH)**

BY

ADEWUMI, GBENGA ADEDEJI

B.Sc. (Hons.) Food Science and Technology (Abeokuta)

M.Sc. Microbiology (Ibadan)

A THESIS SUBMITTED TO THE SCHOOL OF POSTGRADUATE STUDIES,
UNIVERSITY OF LAGOS, NIGERIA FOR THE AWARD OF DOCTOR OF
PHILOSOPHY (Ph.D.) IN MICROBIOLOGY

JANUARY, 2016

CERTIFICATION

DEDICATION

For

Providence Ademidun

Fortune Adebola

Heritage Adedamola

Olufunmilola Rebecca Adewumi (nee Osinowo)

Papa J.I. and Mama C.O. Osinowo

Late Papa J.A. and Mama R.M. Adewumi

ACKNOWLEDGEMENTS

I acknowledge God Almighty for His mercies, loving kindness, guidance and provisions over my life. He has made it possible for me to be alive to witness this day, to Him be all the glory. I am most grateful to my supervisor, Dr. F.A. Oguntinyinbo, who not only initiated this research work, but also took time to guide, mentor and saw it to a successful completion. I equally appreciate Prof. O.O. Omidiji, my co-supervisor, for his advice and guidance during the course of this work. I acknowledge the significant contributions of my postgraduate teachers during the course of this programme; worthy of mention are Professors M.O. Ilori and E.O. Ugoji. I wish to also thank Prof. O.O. Amund for his fatherly interest, and for providing useful materials during my thesis write-up.

The take-off funding for this project was provided by Society for Applied Microbiology (SfAM), Brighton, United Kingdom, through the award of a New Lecturer grant to Dr. F.A. Oguntinyinbo. Government of India provided two postgraduate fellowships, namely DBT-TWAS and INSA JRD-TATA, which enabled my research visits to Institute of Bioresources and Sustainable Development (IBSD), Imphal and National Dairy Research Institute (NDRI), Karnal, India. My utmost appreciation to my hosts, Dr. K. Jeyaram and Dr. S. Grover of IBSD and NDRI respectively, for their special interests in my Ph.D. research. I wish to appreciate my laboratory mates, Wahengbam Romi, Thangjam Anand Singh, Khunjamayum Romapati Devi, Santosh Keisam, Namita Rokana, Hogarehalli Mallapa Rashmi, Rajbir Singh and Dipak Chaudhari, who assisted greatly during my research stay at IBSD and NDRI.

I sincerely acknowledge my employer, University of Lagos, for the award of a doctoral assistance grant, and short-term study leave that facilitated my collaboration with NDRI. At this juncture, I would like to thank the Head, Department of Microbiology, Prof. I.A. Adeleye, for recommending me for this grant. My profound gratitude goes to Bells

University of Technology, Ota, for providing my first teaching employment, and the leave of absence granted during my research visit at IBSD.

To my siblings and numerous friends, especially Dr. C.G. Alimba, I cherish all your counsel and encouragements. Finally, I thank my wife, for her invaluable and outstanding support, my children, for their immeasurable understanding during my long absence from home.

TABLE OF CONTENTS

CONTENTS	PAGES
Title Page	i
Certification	ii
Dedication	iii
Acknowledgements	iv
Table of Contents	vi
List of Tables	xii
List of Figures	xiii
List of Plates	xvi
List of Appendices	xvii
Abstract	xviii
CHAPTER ONE	1
1.0 Introduction	1
1.1 Background to the Study	2
1.2 Statement of the Problem	5
1.3 Aim and Objectives	6
1.3.1 Aim	6
1.3.2 Specific Objectives	6
1.4 Significance of the Study	6
1.5 Operational Definition of Terms	7
CHAPTER TWO	9
2.0 Literature Review	9
2.1 Fermented foods	10
2.2 Fermented vegetable protein seeds of Africa and Asia	10

2.3 Nutritional values of <i>iru</i> and biochemical changes during production	12
2.4 Microorganisms associated with fermented condiments	13
2.5 Microbiological methods for assessment of microbiota in fermented foods	13
2.5.1 Traditional culture-dependent methods	13
2.5.2 Culture-dependent molecular techniques	14
2.5.3 Culture-independent molecular analyses	18
2.6 <i>Bacillus cereus</i> phylogeny: characteristics and identification	25
2.7 Probiotics	32
2.7.1 History and definitions of probiotics	32
2.7.2 Criteria for selecting good probiotic strains	32
2.7.3 Mechanisms of actions of probiotic microorganisms	35
2.7.4 Significant benefits of probiotics	35
CHAPTER THREE	41
3.0 Materials and Methods	41
3.1 Description of samples collection	42
3.2 Bacterial community profiling of <i>iru</i> samples	45
3.2.1 Culture-independent molecular analysis of bacterial community structure in <i>iru</i> samples	45
3.2.1.1 DNA extraction of <i>iru</i> samples	45
3.2.1.2 PCR-DGGE amplification of variable V3 region of 16S rRNA gene	47
3.2.1.3 DGGE analysis of bacterial community of <i>iru</i> samples	48
3.2.1.4 PCR-DGGE gel images analysis	49
3.2.1.5 Excision of DGGE bands and sequencing of amplicons	
3.3 Genomic characterization and sub-typing of autochthonous <i>Bacillus</i> species	50

in *iru*

3.3.1 Culture-dependent microbiological analysis of bacterial species in <i>iru</i> samples	50
3.3.1.1 Isolation of microorganisms	50
3.3.1.2 Determination of Hydrogen ion concentration (pH) of <i>iru</i> samples	51
3.3.1.3 Phenotypic characterization of bacterial species isolated from <i>iru</i> samples	51
3.3.1.3.1 Cultural/colonial/macrosopic characteristics	51
3.3.1.3.2 Gram's staining	51
3.3.1.3.3 Endospore staining	52
3.3.1.3.4 Phase contrast microscopy	52
3.3.1.3.5 Catalase test	52
3.3.1.3.6 Starch hydrolysis	53
3.3.1.3.7 Casein hydrolysis	53
3.3.1.3.8 Lipid hydrolysis	54
3.3.1.3.9 Anaerobic growth	54
3.3.1.3.10 API 50 CHB carbohydrates fermentation	54
profiles and identification system of <i>Bacillus</i> strains	
3.3.2 Culture-dependent molecular assessment of bacterial species in <i>iru</i> samples	55
3.3.2.1 Bacterial genomic DNA extraction, quantification and purity determination	55
3.3.2.2 PCR amplification of 16S rRNA gene	56

3.3.2.3 Amplified ribosomal DNA restriction analysis (ARDRA)	57
3.3.2.4 16S-23S rRNA gene internal transcribed spacer (ITS)	58
PCR amplification	
3.3.2.5 16S-23S rRNA internal transcribed spacer-restriction	59
fragment length polymorphism (ITS-PCR-RFLP)	
3.3.2.6 Randomly amplified polymorphic DNA PCR	59
(RAPD-PCR)	
3.3.2.7 Multilocus sequence analysis (MLSA)	61
3.3.2.8 Phylogenetic construction of 16S rRNA and	61
housekeeping genes sequences	
3.4 Probiotic examinations of <i>Bacillus</i> strains	64
3.4.1 Resistance of <i>Bacillus</i> strains to acidic pH and bile salts tolerance	64
3.4.2 Antimicrobial activity of <i>Bacillus</i> strains against food-borne	64
pathogens	
3.4.3 Bacterial cell surface hydrophobicity (BCSH)	66
3.4.4 Autoaggregation assay	67
3.4.5 Coaggregation assay	67
3.4.6 Cell culture studies	68
3.4.6.1 HT-29 cell lines propagation and maintenance	68
3.4.6.2 <i>In vitro</i> adhesion assay of <i>Bacillus</i> strains	69
3.4.6.3 Bacilli inhibition of <i>S. enterica</i> serovar Typhimurium	70
MBU 1047 adherence to HT-29 cell lines	
3.4.7 Safety assessment of <i>Bacillus</i> strains	72
3.4.7.1 Haemolysis on blood agar	72
3.4.7.2 Antibiotic susceptibility testing	72

3.4.7.3 Mucin degradation by Petri-dish assay	73
3.5 Statistical analysis	74
CHAPTER FOUR	75
4.0 Results	75
4.1 Analysis of bacterial communities and diversity of <i>iru</i> using PCR-DGGE	76
4.2 Identification of major bacterial PCR-DGGE bands	81
4.3 Bacterial counts and pH of <i>iru</i> samples	84
4.4 Phenotypic characterization and identification of bacterial species isolated from <i>iru</i>	87
4.5 <i>Bacillus</i> genomic characterization and strains sub-typing using polyphasic approaches	90
4.6 Phylogenetic analysis	102
4.7 MLSA for sub-typing <i>Bacillus cereus sensu lato</i>	104
4.8 Probiotic functional properties of <i>Bacillus</i> strains	108
4.8.1 Acid resistance and bile salts tolerance	108
4.8.2 <i>In vitro</i> antagonistic potentials of <i>Bacillus</i> strains against food-borne pathogens	110
4.8.3 Bacterial hydrophobicity	113
4.8.4 Autoaggregation	115
4.8.5 Coaggregation	115
4.8.6 Adhesion of <i>Bacillus</i> strains to HT-29 cell lines	117
4.8.7 Inhibitory effects of <i>Bacillus</i> strains on adhesion to HT-29 cell lines by <i>S. enterica</i> serovar Typhimurium MBU 1047	117
4.9 <i>In vitro</i> safety evaluation of <i>Bacillus</i> strains	120
4.9.1 Haemolytic activity	120

4.9.2 Antibiotic susceptibility pattern	120
4.9.3 Degradation of porcine mucin by plate assay	122
CHAPTER FIVE	124
5.0 Discussion	124
5.1 Summary of Findings	151
5.2 Conclusion	152
5.3 Contributions to Knowledge	153
References	154
Appendices	215

LIST OF TABLES

Tables	Pages
2.1 Selection criteria for probiotic strains	34
3.1 Coordinate descriptions of <i>iru</i> sampled at different locations in Nigeria	44
3.2 List of PCR primers for <i>Bacillus</i> species characterization and identification using culture-dependent molecular techniques	63
4.1 Bacterial species richness estimates (<i>R</i>) and Shannon's index of diversity (<i>H</i>) of DGGE profiles of <i>iru</i> samples obtained from various geographical locations in Nigeria	80
4.2 Identities of major bacterial bands excised from PCR-DGGE gels of <i>iru</i> samples	82
4.3 Identities of major bacterial bands excised from PCR-DGGE gels of <i>iru</i> samples	89
4.4 Genomic characterization and sub-typing of <i>Bacillus</i> species isolated from <i>iru</i> , traditional fermented <i>Parkia biglobosa</i> seeds	92
4.5 Antagonistic activities of <i>B. subtilis</i> strains isolated from <i>iru</i> and reference probiotic <i>B. clausii</i> UBBC-07 against eleven indicator bacteria strains	111
4.6 Antibiotic sensitivity profiles of <i>B. subtilis</i> strains isolated from <i>iru</i> and reference probiotic <i>B. clausii</i> UBBC-07	121

LIST OF FIGURES

Figures	Pages
3.1 Map showing sampling locations of <i>iru</i> obtained from different manufacturers and retail markets in South-West and North-Central Nigeria	43
4.1 PCR amplification of variable V3 (~ 240 bp) region of bacterial 16S rRNA gene in different <i>iru</i> samples, based on two genomic DNA extraction protocols- enzymatic (1) and chemical (2)	78
4.2 Principal component of non-metric multidimensional scaling analysis of DGGE data	79
4.3 DGGE profiles of PCR-amplified 16S rRNA gene fragments of sixteen (16) <i>iru</i> samples (alphabets A-P) showing major bacterial amplicons	83
4.4 Population counts of bacterial colonies in <i>iru</i> samples obtained from Oyo (OYO), Abeokuta (ABK), Kaduna (KAD), Ado-Ekiti (ADK), Lagos (LAG), Ibadan (IBA) and Ilorin (ILO)	85
4.5 pH values of <i>iru</i> samples obtained from Oyo (OYO), Abeokuta (ABK), Kaduna (KAD), Ado-Ekiti (ADK), Lagos (LAG), Ibadan (IBA) and Ilorin (ILO)	86
4.6 Phase contrast microscope of <i>Bacillus</i> species endospores	88
4.7 ARDRA gel profiles based on (a) <i>Hae</i> III (b) <i>Cfo</i> I (c) <i>Hinf</i> I (d) <i>Dde</i> I (e) <i>Taq</i> I (f) <i>Rsa</i> I showing differentiation of <i>Bacillus</i> species into phylogeny of <i>B. subtilis</i> and <i>B. cereus</i>	93
4.8 ARDRA gel profile based on <i>Rsa</i> I showing differentiation among <i>B. subtilis</i> phylogeny	94
4.9 ARDRA gel profile based on <i>Cfo</i> I showing differentiation of <i>B. subtilis</i> and <i>B. licheniformis</i>	94

4.10 ITS-PCR gel profile of different <i>B. licheniformis</i> strains	95
4.11 ITS-PCR-RFLP gel profile showing strain diversity within <i>B. subtilis</i> , <i>B. pumilus</i> and <i>B. amyloliquefaciens</i> , also differentiating <i>B. pumilus</i> from <i>B. amyloliquefaciens</i>	95
4.12 Dendrogram based on UPGMA clustering of Jaccard similarity coefficient (S_j) of normalized combined ARDRA, ITS-PCR and ITS-PCR-RFLP fingerprints of <i>Bacillus</i> species isolated from <i>iru</i> and reference strains	96
4.13 RAPD-PCR (OPA 18 primer) of <i>B. subtilis</i> strains from <i>iru</i>	97
4.14 RAPD-PCR (OPA 18 primer) of <i>B. amyloliquefaciens</i> strains from <i>iru</i>	97
4.15 Dendrogram based on UPGMA clustering of Jaccard similarity coefficient (S_j) of normalized OPA 18 RAPD-PCR fingerprints of dominant <i>B. subtilis</i> and <i>B. amyloliquefaciens</i> strains from <i>iru</i> and reference strains	98
4.16 RAPD-PCR (M13 primer) of <i>B. subtilis</i> strains from <i>iru</i>	99
4.17 RAPD-PCR (M13 primer) of <i>B. amyloliquefaciens</i> strains from <i>iru</i>	99
4.18 Dendrogram based on UPGMA clustering of Jaccard similarity coefficient (S_j) of normalized M13 RAPD-PCR fingerprints of dominant <i>B. subtilis</i> strains	100
4.19 Dendrogram based on UPGMA clustering of Jaccard similarity coefficient (S_j) of normalized M13 RAPD-PCR fingerprints of dominant <i>B.</i> <i>amyloliquefaciens</i> strains	101
4.20 Phylogenetic tree showing multiple sequence alignment of 16S rRNA gene sequences of <i>Bacillus</i> species isolated in Africa from <i>iru</i> or <i>daddawa</i> ; <i>hawaijar</i> , <i>cheonggukjang</i> , <i>kinema</i> , <i>douchi</i> , <i>thua nao</i> in Asia; referenced and typed strains	103
4.21 RAPD-PCR (M13 primer) of representative <i>B. cereus</i> strains from <i>iru</i>	105

4.22 PCR amplicons of housekeeping genes (<i>rpoA</i> , <i>gyrB</i> , <i>pheS</i> , <i>glpF</i> and <i>gmk</i>) of <i>B. cereus</i> group from <i>iru</i> based on RAPD-PCR M13 grouping	105
4.23 Phylogenetic relationship of pairwise and multiple alignments of (a) <i>gyrB</i> (b) <i>glpF</i> (c) <i>gmk</i> genes sequences of <i>B. cereus</i> phylotype obtained from <i>iru</i> , and other food and environmental samples, including referenced and typed strains	107
4.24 Growth kinetic and viability of <i>Bacillus</i> strains under simulated gastric juice and intestinal bile conditions	109
4.25 Cell surface adhesion of <i>B. subtilis</i> U170B and <i>B. subtilis</i> U146A from <i>iru</i> and reference probiotic <i>B. clausii</i> UBB-07 to <i>n</i> -hexadecane, toluene and chloroform	114
4.26 Time-course of (a) autoaggregation and (b) coaggregation characteristics of <i>B. subtilis</i> U170B and <i>B. subtilis</i> U146A from <i>iru</i> and reference probiotic <i>B. clausii</i> UBB-07 for a period of 4 h incubation at 37°C	116
4.27 Percentage adhesion of <i>B. subtilis</i> strains U170B, U146A and <i>B. clausii</i> UBB-07 that bound to undifferentiated and differentiated cultured HT-29 cells	119
4.28 Inhibition of adhesion of <i>S. enterica</i> serovar Typhimurium MBU 1047 during exclusion, competition and displacement assays in the presence of <i>B. subtilis</i> strains, U170B, U146A and <i>B. clausii</i> UBB-07	119

LIST OF PLATES

Plates	Pages
4.1 Petri dish containing cell-free extract of <i>B. clausii</i> UBBC-07 showing inhibitory activity against <i>B. cereus</i> MBU 1011 by agar well diffusion method on BHI agar	112
4.2 Mucin degradation assay in agarose Petri dish without glucose	123

LIST OF APPENDICES

Appendices	Pages
I	215
II	216