Oladele et al. Beni-Suef University Journal of Basic and Applied Sciences
(2019) 8:19
https://doi.org/10.1186/s43088-019-0024-x

Beni-Suef University Journal of
Basic and Applied Sciences

RESEARCH **Open Access**

Response of Bambara groundnut (*Vigna subterranean L.*) and Maize (*Zea mays L.*) to heavy metal stress

E. O. Oladele[1*], O. O. Adewumi[1], T. Yahaya[2] and I. A. Taiwo[3]

Abstract

Background: Plants are usually the target of environmental pollution. This study, therefore, investigates the effects of Zinc (Zn) and lead Pb on Bambara nut (*Vigna subterranean*) and Maize (*Zea mays*) at different concentrations, as well as the possible ameliorating effect of the chelant; ethylene diamine acetate (EDTA) and farmyard manure, on the enzymatic activities, the chlorophyll, total protein, and carbohydrate contents.

Results: Findings revealed that Pb and Zn increased the superoxide dismutase (SOD), peroxidase, glutathione synthetase (GSH), malondialdehyde (MDA), and catalase levels with increased concentrations from 100 mg/kg to 200 mg/kg significantly ($p < 0.05$) compared to the control Maize plants. While in Bambara nut, the superoxide dismutase (SOD), peroxidase, glutathione synthetase (GSH), and catalase levels decreased with increased concentrations from 100 mg/kg to 200 mg/kg compared to control except for the malondialdehyde (MDA) which was increased. For Bambara groundnut, with increased Pb concentrations, the chlorophyll content reduced from 2.94 to 2.00 mg/g. However, there was an increase (up to 4.918 mg/g) in the chlorophyll content with increased zinc nitrate concentrations augmented with EDTA at the highest concentration. Maize plants treated with Pb augmented with farmyard manure showed an increase in chlorophyll content with increased concentrations while those assisted with EDTA still experienced a decrease as metal concentrations increased. Bambara groundnut plant had a mean carbohydrate (%) of 14.79 (control), 17.60 (100 mg/kg of Pb concentration) and 11.20 (200 mg/kg of Pb concentration), indicating a decrease in carbohydrate content with increased Pb concentrations. The same trend was observed for the different Zn and Pb concentrations on the mean total proteins and carbohydrates of both test plants. Generally, Pb and Zn induced oxidative stress in treated plants.

Conclusions: Elevated activity of anti-oxidative enzymes can assist as important components of antioxidative defense mechanism against oxidative damage. The results of this study could be beneficial in the understanding of the role of the defense system as well as the detoxification mechanism of *Vigna subterranean* and *Zea mays* in efficient tolerance and response to Pb and Zn. This signifies that these plants can act as bioindicators in environmental quality assessment.

Keywords: Heavy metals, Superoxide dismutase (SOD), Phytochelatins, Glutathione synthetase (GSH), Malondialdehyde (MDA)

* Correspondence: eoladele@unilag.edu.ng
[1]Biology Unit, Distance Learning Institute, University of Lagos, Akoka, Lagos, Nigeria
Full list of author information is available at the end of the article

Springer Open

© The Author(s). 2019 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

2.2 Experimental description

The soil was mixed thoroughly and then filled into 60 black cellophane bags. Four thousand grams (4 kg) of soil were placed in each bag. The bags were arranged in four rows designated as control (untreated soil) and soil with metals. The experiment was carried out under a period of 90 days [48].

2.3 Heavy metal determination in the plant samples

The plant samples were digested with concentrated HNO_3 + $HClO_4$ following a modified method described by Lone et al. (2008). The samples were analyzed for lead (Pb) and zinc (Zn) accumulation (mg kg^{-1} DW) using Atomic Absorption Spectrometer (AAS) [28].

2.4 Determination of superoxide dismutase SOD activity

The level of determination of superoxide dismutase (SOD) activity in the control and treated plant samples were determined by the method of Magwere et al. [30]. Superoxide dismutase was assayed by monitoring the inhibition of photochemical reaction into nitro blue tetrazolium (NBT). A unit of SOD activity is expressed as the amount of enzyme required to cause 50% inhibition in the reduction of nitro blue tetrazolium (NBT) at 560 nm.

2.5 Determination of GSH activity

The total sulphydryl groups, protein-bound sulphydryl groups, and free sulphydryl groups like glutathione (GSH) in biological samples was determined using Ellman's reagent, 5,5′-dithio-bis-2-nitrobenzoic acid (DTNB) according to the modified method by Jollow et al [26]. Fresh leaf tissue (1 g) was ground in potassium phosphate buffer 950 Mm; pH 7) containing EDTA (1 mM) and insoluble PVP (2%). The homogenate was centrifuged for 10 min at 15,000 g and the supernatant was used as enzyme extract.

2.6 Determination of CAT activity

Catalase (CAT) activity of the control and treated plant samples were determined according to the method of Sinha [40] but modified by Artenie et al. [8]. Different amounts of H_2O_2 ranging from 20 to 160 moles were taken in small test tubes and 2 ml of dichromate/acetic acid was added to each. The addition of the reagents instantaneously produces an unstable blue precipitate of perchromic acid. Subsequent heating for 10 min in a boiling water bath changed the color of the solution to stable green due to the formation of chromic acetate. After cooling at room temperature, the volume of the reaction mixture was made to 3 ml with distilled water and the absorbance measured with a spectrophotometer at 570 nm.

2.7 Determination of product of lipid peroxidation (MDA)

This was assayed by measuring the TBA (thiobarbituric acid), reactive products present in the treated and untreated plant samples using the procedure of Vashney and Kale [44] and expressed as micromolar of malondialdehyde (MDA)/g tissue. Lipid peroxidation was measured in terms of malondialdehyde (MDA) content. An aliquot of 0.4ml of the test sample was mixed with 1.6 ml of Tris KCl buffer (which was first placed into the test tube before the test sample). Then, 0.5 ml of 30% TCA (Trichloroacetic acid) was added followed by 0.5 ml of 0.75% TBA (Thiobarbituric acid) and the mixture was placed in a water bath for 1hour between 90–95 °C. This was then cooled in ice and centrifuged at 3000 rpm. for 15 min. The clear pink supernatant was collected and absorbance measured against a reference blank of distilled water at 532 nm in a spectrophotometer.

2.8 Determination of chlorophyll content in treated and untreated plant samples

The leaf total chlorophyll content (%) was evaluated. The chlorophyll content of the seedling was determined using the method of Arnon [7]. Leaves from plants from each treatment and the control were separately put in a clean mortar, 10 ml of 80% acetone was added and the leaf tissue was ground to fine pulp for 3 min. The resulting green liquid was carefully transferred into a Buchner funnel containing a pad of Whatman No. 1 filter paper. The optical density (OD) of the chlorophyll extract was determined with a spectrophotometer at 663 nm and 645 nm against a 100% acetone solvent blank. (Perkin Elmer UV/VIS Lambda Bio Spectrophotometer)

2.9 Proximate analysis

The following parameters were determined for the proximate analysis using the leaves:

Total carbohydrates were determined, after hydrolysis, colorimetrically using anthrone reagent (Fales, [20]). Total carbohydrate contents were measured using the phenol-sulfuric acid assay and using glucose as a standard following the method described by DuBois et al. [18].

Total protein was obtained by determining the organic nitrogen content of the sample using the method by Gopal and Rizvi [22] and multiplying the % nitrogen by a protein conversion factor which is usually 6.25 [27]. Using the method described by Gopal and Rizvi [22], total protein was colorimetrically measured at 540 nm against blank using spectrophotometer.

2.10 Statistical analysis

All data collected were analyzed using standard deviation, t test, and analysis of variance (ANOVA) for

statistical significance at 95% confidence interval. Descriptive statistics were calculated using the Microcal origin 5.0 and Microsoft Excel. Graphical illustrations were also carried out to get a vivid representation of the data obtained.

3 Results

Table 1 shows the effects of lead and zinc treatments on superoxide dismutase (SOD), peroxidase, glutathione (GSH), catalase activity, and the level of malondialdehyde (MDA) in Maize. Zn and Pb treatment significantly increased the activities of CAT, peroxidase enzymes, and the level of MDA compared to control ($P < 0.05$) in Maize. With increased zinc concentration, the GSH activity increased with a significant difference from the untreated plants ($P < 0.05$). Also, at the highest Pb concentration of 200 mg/kg, the GSH activity reduced with a significant difference compared to the untreated plants ($P < 0.05$). The SOD activity was also raised with increased zinc concentration with significant difference ($P < 0.05$) compared to control. However, at the lowest Pb and Zn concentrations of 100 mg/kg, there was no significant difference ($P > 0.05$) in the SOD level compared to control. At the highest lead concentration, the SOD level observed was significantly higher than the control ($P < 0.05$).

Table 2 shows the effect of lead and zinc treatment on superoxide dismutase (SOD), peroxidase, glutathione (GSH), catalase activities, and the level of malondialdehyde (MDA) in Bambara groundnut. Zn and Pb treatment significantly decreased enzyme activity compared to control ($P < 0.05$) except for the MDA in Bambara nut treated with 100 mg/kg zinc nitrate. The SOD and catalase activities were significantly reduced, while the MDA increased.

3.1 Chlorophyll content

Table 3 shows the effects of the different concentrations of the metals on the total chlorophyll content of treated plants. The chlorophyll content (mg/g) of treated plants decreased significantly with increased lead and zinc concentrations in all treated plants. The chlorophyll contents of all treated plants were affected significantly at all concentrations. The higher concentration of metals decreased the quantity of total chlorophyll of plants significantly ($P < 0.05$) when compared to the untreated (control) plants. For Bambara groundnut, with increased Pb concentrations the chlorophyll content reduced. However, there was an increase in the chlorophyll content with increased zinc nitrate concentrations augmented with EDTA. Maize plants treated with Pb augmented with farmyard manure showed an increase in chlorophyll content as concentrations of Pb increased while those assisted with EDTA still experienced a decrease rather than an increase in chlorophyll contents as the metal concentrations increased. The untreated (control) Bambara groundnut plant had mean total chlorophyll (mg/g) of 1.598 ± 0.001, and Bambara groundnut treated with 100 mg/kg of lead nitrate had a mean total chlorophyll (mg/g) of 2.944 ± 0.952 and the plant treated with 200 mg/kg of lead concentration had mean total chlorophyll (mg/g) of 2.002 ± 0.616. Bambara groundnut plants with 200 mg/kg of Pb augmented with EDTA gave mean chlorophyll content (mg/g) of 2.193 ± 0.117, while the 200 mg/kg of Pb augmented with manure gave 5.418 ± 0.145. Maize control plants had a mean chlorophyll content of 2.350 ± 0.03 (mg/g). Maize treated with 100 mg/kg and 200 mg/kg of Pb concentration had mean chlorophyll content (mg/g) of 1.242 ± 0.02 and 1.092 ± 0.01, respectively. Maize plants treated with 200 mg/kg of Pb augmented with EDTA gave mean chlorophyll content (mg/g) of 1.876 ± 0.02, while the plant treated with 200 mg/kg of Pb augmented with manure gave 1.621 ± 0.04 (mg/g). Maize plants treated with Zn augmented with farmyard manure showed an increase in chlorophyll content as concentrations of Zn increased while those assisted with EDTA still experienced a decrease rather than an increase in chlorophyll contents as the metal concentrations increased. The untreated (control) Bambara groundnut plant had mean total chlorophyll (mg/g) of 1.598 ± 0.001, Bambara groundnut treated with

Table 1 Impact of different concentrations of Lead and Zinc nitrates on some Enzymes and Lipid Peroxidation in maize

Treatment	SOD (U/mg)	PEROXIDASE (U/mg)	GSH (μmole/min/mg)	CAT (U/mg)	MDA (U/mg)
Control	0.75 ± 0.03	18.20 ± 0.15	0.26 ± 0.08	16.54 ± 0.44	0.008 ± 0.002
100 mg/kg lead	0.78 ± 0.02	17.58 ± 0.10*	0.08 ± 0.03*	24.43 ± 0.37*	0.118 ± 0.018*
150 mg/kg lead	0.98 ± 0.22*	22.20 ± 0.27*	0.12 ± 0.02*	27.54 ± 0.37*	0.310 ± 0.02
200 mg/kg lead	1.10 ± 0.11*	26.10 ± 0.50*	0.17 ± 0.02*	30.42 ± 0.07*	0.430 ± 0.04*
100 mg/kg zinc	0.82 ± 0.06	23.40 ± 0.25*	0.18 ± 0.04*	22.21 ± 0.30*	0.080 ± 0.02*
150 mg/kg zinc	2.77 ± 0.24*	46.40 ± 0.15*	0.21 ± 0.07*	72.10 ± 0.12*	0.202 ± 0.005*
200 mg/kg zinc	2.98 ± 0.45*	52.20 ± 0.44*	0.30 ± 0.03*	76.22 ± 0.15*	0.345 ± 0.015*

The values are the means ± SEM (range) of 3 replicates
When *$P < 0.05$ = significantly different from control
When $P > 0.05$ = not significantly different from control

Table 2 Impact of different concentrations of lead and zinc nitrate on some enzymes and lipid peroxidation in Bambara nut

Treatment	SOD (U/mg)	PEROXIDASE (U/mg)	GSH (μmole/min/mg)	CAT (U/mg)	MDA (U/mg)
Control	2.57 ± 0.05	53.42 ± 0.83	0.45 ± 0.03	72.42 ± 0.39	0.033 ± 0.007
100 mg/kg lead	1.00 ± 0.03*	38.48 ± 0.53*	0.12 ± 0.03*	40.08 ± 0.24*	0.068 ± 0.005*
150 mg/kg lead	0.72 ± 0.02*	20.86 ± 0.14*	0.09 ± 0.02*	20.35 ± 0.35*	0.071 ± 0.002*
200 mg/kg lead	0.68 ± 0.05*	20.10 ± 0.14*	0.06 ± 0.01*	18.65 ± 0.35*	0.092 ± 0.002*
100 mg/kg zinc	1.86 ± 0.05*	51.44 ± 0.05*	0.41 ± 0.02*	66.78 ± 0.12*	0.029 ± 0.011
150 mg/kg zinc	1.69 ± 0.05*	48.40 ± 0.55*	0.38 ± 0.02*	64.38 ± 0.29*	0.038 ± 0.002*
200 mg/kg zinc	1.22 ± 0.04*	44.30 ± 0.40*	0.29 ± 0.03*	54.34 ± 0.09*	0.082 ± 0.002*

The values are the means ± SEM (range) of 3 replicates
When *$P < 0.05$ = significantly different from control
When $P > 0.05$ = not significantly different from control

100 mg/kg of zinc nitrate had a mean total chlorophyll (mg/g) of 3.649 ± 0.13 and the plant treated with 200 mg/kg of Zn concentration had mean total chlorophyll (mg/g) of 3.112 ± 0.06. Bambara groundnut plants with 200 mg/kg of Zn augmented with EDTA gave mean chlorophyll content (mg/g) of 4.918 ± 0.21, while the 200 mg/kg of Zn augmented with manure gave 5.816 ± 0.16. Maize control plants had a mean chlorophyll content of 2.350 ± 0.03 (mg/g). Maize treated with 100 mg/kg and 200 mg/kg of Zn concentration had mean chlorophyll content (mg/g) of 1.565 ± 0.04 and 1.352 ± 0.03, respectively. Maize plants treated with 200 mg/kg of Zn augmented with EDTA gave mean chlorophyll content (mg/g) of 3.989±0.07, while the plant treated with 200mg/kg of Zn augmented with manure gave 6.101±0.15 (mg/g).

3.2 Proximate analysis

3.2.1 Total carbohydrate and total protein

Tables 4 and 5 show the effects of different Lead and Zinc concentrations on the total protein and total carbohydrates. The total protein and total carbohydrate of treated plants decreased significantly with increased lead and zinc concentrations. For Maize, whether assisted with chelator and manure or not, the protein and carbohydrate contents decreased significantly with increased lead and zinc concentrations. However, farmyard manure assisted Bambara groundnut when planted in soil with the lowest lead concentrations of 100 mg/kg and different zinc nitrate concentrations. They had higher percentages of protein and carbohydrates compared with the mean values observed for their control plants. When the Bambara groundnut was planted in soil containing the lowest concentration of lead and mixed with EDTA, a high-percentage carbohydrate was also observed. Control Bambara groundnut plant had a mean carbohydrate (%) of 14.79 ± 0.09, 17.60 ± 0.01 for 100 mg/kg of lead concentration and 11.20 ± 0.02 for 200 mg/kg of lead concentration, indicating a decrease in carbohydrate content with increased concentrations. Bambara groundnut plants with 100 mg/kg of lead and EDTA gave a mean percentage carbohydrate of 15.57 ± 0.08,

Table 3 Impact of lead concentrations on total chlorophyll (mg/g) of the test plants

Plants/concentration	Bambara groundnut (Pb treatment) Mean value ± SEM	Maize (Pb treatment) Mean value ± SEM	Bambara groundnut (Zn treatment) Mean value ± SEM	Maize (Zn treatment) Mean value ± SEM
Control	1.598 ± 0.001	2.350 ± 0.03	1.598 ± 0.001	2.350 ± 0.03
100 mg/kg	2.944 ± 0.952*	1.242 ± 0.02*	3.649 ± 0.13*	1.565 ± 0.04*
150 mg/kg	2.418 ± 0.202*	1.108 ± 0.02*	3.501 ± 0.1*	1.486 ± 0.05*
200 mg/kg	2.002 ± 1.616*	1.092 ± 0.01*	3.112 ± 0.06*	1.352 ± 0.03*
100 mg/kg + EDTA	2.872 ± 0.618*	2.028 ± 0.03*	3.712 ± 0.09*	4.141 ± 0.10*
150 mg/kg + EDTA	2.648 ± 0.320*	1.912 ± 0.12*	4.518 ± 0.10*	4.102 ± 0.11*
200 mg/kg + EDTA	2.193 ± 0.117*	1.876 ± 0.02*	4.918 ± 0.21*	3.989 ± 0.07*
100 mg/kg + manure	5.712 ± 0.100*	1.308 ± 0.01*	4.612 ± 0.09*	5.122 ± 0.11*
150 mg/kg + manure	5.528 ± 0.150*	1.475 ± 0.03*	5.622 ± 0.18*	5.893 ± 0.06*
200 mg/kg + manure	5.418 ± 0.145*	1.621 ± 0.04*	5.816 ± 0.16*	6.101 ± 0.15*

The values are the means ± SEM (range) of 3 replicates
When *$P < 0.05$ = significantly different from control
When $P > 0.05$ = not significantly different from control

Table 4 Effect of lead and zinc concentration on the total carbohydrate of the test plants

Plants/concentration	Bambara groundnut	Maize	Bambara groundnut	Maize
	Lead treatment		Zinc treatment	
	Mean value ± SEM	Mean value ± SEM	Mean value ± SEM	Mean value ± SEM
Control	14.79 ± 0.44	17.26 ± 0.94	14.79 ± 0.09	17.26 ± 0.14
100 mg/kg	17.60 ± 0.01*	17.30 ± 0.08*	6.75 ± 0.2*	11.33 ± 0.03*
150 mg/kg	12.10 ± 0.02*	14.20 ± 0.01*	5.98 ± 0.02*	9.29 ± 0.04*
200 mg/kg	11.20 ± 0.02*	12.82 ± 0.04*	5.11 ± 0.07*	8.61 ± 0.05*
100 mg/kg + EDTA	15.57 ± 0.08*	9.17 ± 0.03*	13.81 ± 0.11*	12.86 ± 0.11*
150 mg/kg + EDTA	14.21 ± 0.23*	8.64 ± 0.01*	10.42 ± 0.13*	9.54 ± 0.05*
200 mg/kg + EDTA	12.46 ± 0.02*	7.62 ± 0.01*	8.49 ± 0.06*	8.47 ± 0.04*
100 mg/kg + manure	17.58 ± 0.06*	16.91 ± 0.05*	17.81 ± 0.05*	16.42 ± 0.2*
150 mg/kg + manure	8.43 ± 0.05*	7.05 ± 0.21*	14.23 ± 0.02*	7.22 ± 0.05*
200 mg/kg + manure	7.36 ± 0.03*	6.78 ± 0.04*	9.42 ± 0.04*	6.89 ± 0.08*

The values are the means ± SEM (range) of 3 replicates
When *$P < 0.05$ = significantly different from control
When $P > 0.05$ = not significantly different from control

while the 100 mg/kg of lead and Manure gave 17.58 ± 0.06. Maize control plants had 17.26 ± 0.94 mean percentage carbohydrate. Maize treated with 100 mg/kg and 200 mg/kg of lead concentration had a mean percentage carbohydrate of 17.30 ± 0.08 and 12.82 ± 0.04, respectively. Maize plants with 200 mg/kg of lead and EDTA gave a mean percentage carbohydrate of 9.17 ± 0.03, while the 200 mg/kg of lead and manure gave 16.91 ± 0.05. The same trend was observed for the different zinc and lead concentrations on the mean total proteins of the two test plants. The statistical analysis, therefore, shows that the percentage of carbohydrates and proteins of treated plants were significantly different from their control ($p < 0.05$) for each tested crop plant.

4 Discussion

4.1 Biochemical analysis

Plant food generally contains abundant natural bioactive compounds that have health-promoting activities like antihypertensive, anti-inflammatory, and antioxidants, such bioactive compounds therefore include polyphenols [5]. However, when plants' biochemical activities are disrupted due to heavy metal exposure such benefits from the food (protein and carbohydrates) are therefore lost or reduced in plants. In the case of the study plants, the enzymatic activities of Bambara groundnut and Maize were adversely affected when the plants were exposed to different concentrations of lead and zinc. This was probably attributed to oxidative stress induced by these metals. For both test plants, there were increase and

Table 5 Effect of lead and zinc concentration on total protein of the test plants

Plants/concentration	Bambara groundnut	Maize	Bambara groundnut	Maize
	Lead treatment		Zinc treatment	
	Mean value ± SEM	Mean value ± SEM	Mean value ± SEM	Mean value ± SEM
Control	9.52 ± 0.09	6.46 ± 1.0	9.52 ± 1.95	6.46 ± 1.26
100 mg/kg	8.80 ± 0.38*	5.89 ± 1.05*	9.60 ± 0.40*	5.16 ± 0.55*
150 mg/kg	8.40 ± 0.36*	5.63 ± 1.0*	9.10 ± 0.59*	5.04 ± 0.59*
200 mg/kg	7.40 ± 0.30*	5.08 ± 0.64*	8.60 ± 1.03*	4.96 ± 0.13*
100 mg/kg + EDTA	10.09 ± 0.25*	4.04 ± 1.44*	10.12 ± 0.84*	4.18 ± 0.51*
150 mg/kg + EDTA	9.92 ± 0.19*	3.84 ± 0.35*	9.84 ± 1.01*	4.12 ± 0.45*
200 mg/kg + EDTA	9.80 ± 0.16*	3.78 ± 1.10*	9.72 ± 1.47*	3.98 ± 0.24*
100 mg/kg + manure	10.28 ± 0.30*	4.12 ± 0.51*	9.80 ± 1.45*	5.56 ± 0.32*
150 mg/kg + manure	9.68 ± 0.25*	4.20 ± 0.51*	9.64 ± 0.75*	5.97 ± 0.31*
200 mg/kg + manure	8.10 ± 0.20*	4.80 ± 0.95*	9.59 ± 1.53*	6.12 ± 0.54*

The values are the means ± SEM (range) of 3 replicates
When *$P < 0.05$ = significantly different from control
When $P > 0.05$ = not significantly different from control

decrease in the level of the enzymes. The increased level of MDA (malondialdehyde) is highly indicative of oxidative stress for the plants. An increase in the level of MDA (product of lipid peroxidation) can lead to inactivation of enzymes, DNA damage, and interaction with vital plant cells. These results agreed with the work of Mansoor and Nasir [32] and Vazquez et al. [45]. The levels of glutathione, peroxidase, SOD, and catalase for the treated plants were also raised. The increased level of glutathione, peroxidase, and catalase also signifies the possible release of free radicals. This was again observed by Mansoor and Nasir [32]. The level of superoxide dismutase, catalase, peroxidase, and glutathione were lowered in Bambara groundnut, except for the MDA. The low level of superoxide dismutase signifies the possible release of radicals within the treated Bambara groundnut due to no inhibition of oxidation. Malar et al. [31] also reported that the activity of antioxidative enzymes, such as APX and POX, in water hyacinths (*Eichhornia crassipes* (*Mart.*), positively correlated with Pb treatment, while in the case of SOD and CAT enzymes increased up to 800 mg/L treatment and then slightly decreased at higher concentrations. Anee et al. [6] had a similar observation with fluctuating enzyme activities especially, decreased catalase activity when *Sesamum indicum* was subjected to the stress of different waterlogging durations. This indicates that stress levels increased with increased waterlogging.

Also, this could be due to low metal translocation potential displayed by the plants. It could also be suggested as a tolerance capacity displayed by Bambara nut to protect itself from oxidative damage. On the other hand, the lowered MDA content indicates toxicity/poisoning in the test plant. MDA is the end product of peroxidation of membrane lipids and it accumulates when the plants are subjected to oxidative stress [44]. Generally, free radical generation and membrane damage would be low in tolerant plants and thereby formation of lower levels of MDA content [39]. As a result, in this study, the comparatively lowered MDA content in test plants to Pb and Zn stress may assist its tolerant nature. These findings corroborated the earlier work by Dietz and Schnoor [17] that lead stress causes multiple direct and indirect effects on plant growth and metabolism and also alter some physiological processes. Edwards et al. [19] also observed that high internal levels of glutathione can be beneficial as a first response towards some of the effects exerted by pollutants. Cells are usually protected from reactive oxygen species by the combined action of enzymatic antioxidant systems like catalase, peroxidase, and non-enzymatic antioxidant like ascorbate, glutathione, and phenolic compounds [13, 19]. Glutathione (GSH), a precursor for phytochelatins (PCs) which are heavy-metal binding peptides, is known to be involved in heavy-metal tolerance and sequestration. GSH plays a central role in defense against oxidative stress and heavy metals. Lead compounds have been observed to bind less strongly to phytochelatins (PCS) unlike other metals like zinc due to larger ion radius (Pb, octahedral) and high coordination number (Pb, 6-8). This could be the reason for the greater Pb tolerance and uptake by Maize with or without augmentation and less Pb tolerance by Bambara groundnut. This could also be attributed to the great tolerance of Zn exhibited by the two plants.

4.2 Chlorophyll content

The chlorophyll content of treated plants reduced significantly with increased metal concentration especially for lead in all treated plants. As the concentration of metals increased, the quantity of total chlorophyll in the treated plants decreased. This led to a reduction in photosynthetic activities and an indication of stress. This finding therefore agrees with Chettri et al. [16] who reported a decrease in chlorophyll levels for *Cladonia rangiformis* after treating with Cu, Zn, and Pb. They added that a decrease of chlorophyll after Cu application may be due to the blocking of enzymes acting in chlorophyll synthesis or to degradation of chlorophyll. They also said that chlorophyll content is usually measured in plants in order to assess the impact of environmental stress, as changes in pigment content are linked to visual symptoms of plant illness and photosynthetic productivity. The chlorophyll content reduced with increased zinc and lead nitrate concentrations [39]., had similar observations as experienced in this present study, with a reduction in chlorophyll and carotenoid content of cuttings and seedlings of *Jatropha curcas* L. under high concentration of lead (Pb). Shu et al. [39] perceive chlorophyll degradation and photosynthesis inhibition as a specific response of plants to metal stress. The lowered content of chlorophyll specifies a decrease in net photosynthesis due to reduced absorption of essential mineral nutrients as an indirect reason for plant chlorosis. Lead (Pb) contamination induces leaf chlorosis and may eventually lead to cell death [37].

4.3 Total carbohydrate and total protein

The total protein and total carbohydrate of treated plants decreased significantly with increased lead and Zinc concentrations. However, farmyard manure assisted Bambara groundnut when planted in soil with manure mixed with the lowest lead concentrations of 100 mg/kg and different zinc nitrate concentrations. In support of these findings, several studies show that lead (Pb) significantly affects nutrient uptake and overall distribution of nutritional elements within plant organs [9, 23, 49]. Chettri et al. [15], observed that lead (Pb) had an adverse effect on the metabolites content through protein in